# MUS. COMIOCCASIONAL Papers On Mollusks

110V 2 5 1974

Published by

THE DEPARTMENT OF MOLLUSKS

HARVA Museum of Comparative Zoölogy, Harvard University
UNIVERSITY
Cambridge, Massachusetts

VOLUME 4

15 NOVEMBER 1974

NUMBER 49

## CATALOGUE OF THE TAXA OF LUCIDELLA (PROSOBRANCHIA: HELICINIDAE)

By KENNETH J. BOSS AND MORRIS K. JACOBSON

This catalogue was initiated during the preparation of our taxonomic revision of the species of the genus Lucidella which inhabit the island of Cuba (see preceding paper, Boss and Jacobson, 1974, Occ. Papers Moll., no. 48). So much information was collected on nominal species of Lucidella that we thought it was worthwhile to make it available. The catalogue includes: 1) generic, subgeneric, specific and subspecific nomina as originally introduced; 2) the original citation with date; 3) the type-locality; 4) and the whereabouts of type-specimens, if found. In some cases we have also provided references of later date if they present a figure of the taxon.

### ACKNOWLEDGMENTS

We are indebted to the following persons for providing information about type-specimens: Drs. R. Robertson and G. Davis, Academy of Natural Sciences of Philadephia; Dr. J. Rosewater, United States National Museum of Natural History, Washington; Dr. J. J. Oberling, Naturhistorisches Museum, Bern, Switzerland; Dr. A. Riedel, Instytut Zoologiczny, Warszawa, Poland; Dr. A. Zilch, Senckenbergische Gesellschaft, Frankfort, West Germany;

Miss S. M. Wells, British Museum (Natural History), London. Dr. R. D. Turner and Mr. R. I. Johnson reviewed the manuscript.

### ABBREVIATIONS

ANSP — Academy of Natural Sciences, Philadelphia

BMNH — British Museum (Natural History), London

IZW — Instytut Zoologiczny Warszawa, Poland

MCZ — Museum of Comparative Zoology, Cambridge, Massachusetts

NMB — Naturhistorisches Museum, Bern, Switzerland

USNM — United States National Museum of Natural History, Washington

SMF — Senckenbergische Naturforschende Gesellschaft, Frankfort, West Germany.

abbotti Vanatta, Helicina, 1920. Proc. Acad. Nat. Sci. Phila. 72: 205, pl. 6, fig. 12, 14 (1 mi NW of Sanchez, Santo Domingo, Hayti Id. [sic], [Hispaniola]; type, ANSP 44370a).

adamsiana Pfeiffer, Helicina, 1848. Proc. Zool. Soc. London, p. 119 (Jamaica; type, destroyed; Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 344, pl. 68, figs. 18–20). NOTE: type-species of Urichia Guppy 1895.

areola (Férussac) Petit de la Saussaye, Helix, 1851. Jour. de Conch. 2: 86, pl. 1, fig. 10, error for aureola Férussac.

aureola Férussac, Helix, 1822. Histoire naturelle général et particulière des mollusques terrestres et fluviatiles, cover of livraison 17 (see: Kennard, 1942), pl. 48, fig. 1; assigned to the subgenus Helicodonta in Tableaux systématiques des animaux mollusques . . . Prodrome Général . . . Tableau systématique de la famille des Limaçons, no. 90, p. 33 (La Martinique).

barbadensis Pfeiffer, Helicina, 1853. Proc. Zool. Soc. London, p. 60 (Barbadoes; type, destroyed).

beatensis Bartsch, Lucidella, 1932. Proc. U.S. Nat. Mus.

- 81 (2929): 3, pl. 2, figs. 4-6 (Beata Is., Island of Haiti [Dominican Republic, Hispaniola]; type, USNM 403920).
- caymanensis Pilsbry, Lucidella, 1930. Proc. Acad. Nat. Sci. Phila. 82: 230, pl. 15, figs. 1, 1a, 1b (½ mi S of North Side Area, Grand Cayman Id.; type, ANSP 150867).
- christophori Pilsbry, Helicina plicatula, 1897. Nautilus 10: 118 (St. Kitts; type, ANSP 62062a). NOTE: type-species of *Poeniella* H. B. Baker 1923.
- cibaoensis Crosse, Helicina rugosa var., 1891. Jour. de Conch. 39: 185 (Cibao, Santo Domingo, Hispaniola; type, lost); Pilsbry 1928. Proc. Acad. Nat. Sci. Phila. 80: 480 (neotype designated, ANSP 14796a, pl. 27, fig. 1 and 2; Puerto Plata, Santo Domingo, Hispaniola). NOTE: Crosse merely gave a name to var. B of Hjalmarson and Pfeiffer 1858, Malak. Blätt. 5: 145, var. of rugosa.
- conoidea Pilsbry, Lucidella (Poeniella), 1933. Proc. Acad. Nat. Sci. Phila. 85: 133, pl. 7, figs. 7, 7a (Del Monte Plantation, near Salvation, Santo Domingo, 3000 ft.; type, ANSP 160982).
- coronula (Shuttleworth) Pfeiffer, Helicina, 1862. Malak. Blätt. 9: 155 (Jamaica; type, destroyed); Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 345, pl. 68, fig. 21.
- costata Simpson, Lucidella, 1894. Proc. U.S. Nat. Mus. 17: 449, pl. 16, fig. 6 (marl stratum of Miocene beds, Bowden, St. Thomas, Jamaica; type, USNM 115766).
- cruciata Weinland, Helicina, 1880. Jahrb. Malak. Gesell. 7: 351, pl. 12, fig. 7 (Port au Prince, Haiti, Hispaniola; type, not located). NOTE: synonym of paivana Pfeiffer 1866.
- decussata Boettger, Helicina, 1887. Jahrb. Malak. Gesell. 14: 102, pl. 4, fig. 8 (Miragoane, Haiti, [Hispaniola]; type, SMF 181169 a). NOTE: subspecies of paivana Pfeiffer 1866.
- denseplicata Wagner, Lucidella, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 348, pl. 69, figs. 14–15 (St. Lucia, Lesser Antilles; syntypes, IZW 9100).

depressa Gray Helicina, 1825. Zoological Jour. 1: 69, pl. 6, fig. 14 (West Indies; holotype, BMNH 1874.99.8). NOTE: Type-species of Poenia H. and A. Adams 1856.

euglypta Crosse, Helicina, 1874. Jour. de Conch. 22: 119, 204, pl. 4, fig. 4 (near Fort-de-France, Martinique; type, not located). NOTE: Wagner (1910) cited this as a synonym of plicatula Pfeiffer 1848.

foxi Pilsbry, Lucidella, 1899. Nautilus 13: 56 (Cave at Port Antonio, Jamaica; type, ANSP 61848a); Wagner, 1910, [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 343, pl. 68, figs. 10-11.

gonavensis Pilsbry, Lucidella (Poeniella), 1928. Proc. Acad. Nat. Sci. Phila. 80: 481, pl. 27, figs. 8-9 (Gonave Id.,

Haiti, Hispaniola; type, ANSP 146698).

granulosa C. B. Adams, Lucidella aureola, 1850. Contrib. Conch. no. 6: 89 (Jamaica; lectotype, MCZ 156170, Jacobson and Boss, 1973, p. 358, pl. 84, figs. 1-3).

granulum (Gundlach) Pfeiffer, Helicina, 1864. Malak. Blätt. 11: 161 (Monte Toro, district of Guantánamo, Oriente, [Cuba]; type, destroyed; neotype, ANSP 110769, designated and figured by Jacobson, 1973, p. 108, fig. 1).

granum Pfeiffer, Helicina, 1856. Novit. Conch. 1: 86, pl. 23, figs. 20-23; 1857. Malak. Blätt. 3: 49 (Holguín, Oriente, Cuba; type, destroyed; neotype, MCZ 73863, designated and figured, Boss and Jacobson, 1974, p. 11, pl. 1, figs. 1-3).

holoserica Wagner, Lucidella, 1910. [in] Martini and Chemnitz. Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 350, pl. 69, figs. 16-19 (Barbadoes; syntypes, IZW 9101).

ignicoma Guppy, Helicina, 1868. Ann. Mag. Nat. Hist. (4) 1: 441 (Trinidad; type, not in the BMNH, probably lost in the fire of the Victoria Institute, Trinidad, in 1920).

inaequalis Pfeiffer, Helicina (Lucidella), 1859. Proc. Zool. Soc. London, p. 28 (Jamaica; holotype, BMNH 1974.1).

indecora Pilsbry, Lucidella pilsbryi, 1930. Proc. Acad. Nat. Sci. Phila. 82: 243, pl. 17, fig. 5 (Little Swan Id.; [Caribbean Seal; type, ANSP 150881).

interrupta Simpson, Lucidella aureola var., 1894. Proc. U.S. Nat. Mus. 17: 449 (Duncan's, Trelawney, Jamaica; type,

- USNM). NOTE: Probable synonym of montegoensis Brown, fide, H. B. Baker (1934).
- jeremiensis Pilsbry, Lucidella, 1929. Proc. Acad. Nat. Sci Phila. 28: 479, pl. 27, figs. 3-5 (near General's Cave, 1 mi W of Jérémie, Haiti, [Hispaniola]; holotype ANSP 118250a).
- kobelti Wagner, Lucidella, 1910. [in] Martini and Chemnitz Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 350, pl. 69, figs. 12 and 13 (Jamaica; holotype, IZW 9097).
- lamellosa Guppy, Helicina (Perenna), 1867. Ann. Mag. Nat. Hist. (3) 19: 260, pl. 10, fig. 4 (Trinidad; syntype, BMNH 1874.10.30.12). NOTE: synonym of H. lirata Pfeiffer 1847; monotype of Perenna Guppy 1859.
- lasellensis Clench and Aguayo, Lucidella (Poeniella), 1937. Mem. Soc. Cubana Hist. Nat. 11 (2): 63, pl. 7, fig. 6 (La Visite, La Selle Range, Haiti, [Hispaniola]; holotype, MCZ 110626).
- leana C. B. Adams, Helicina, 1845. Proc. Boston Soc. Nat. Hist. 2: 11 (Jamaica; type, lost, see Jacobson and Boss, 1973, p. 375). NOTE: C. B. Adams (1851) called this a synonym of depressa Gray.
- Lindsleya Chitty, 1857. Proc. Zool. Soc. London, pp. 167 and 169 (type-species, Stoastoma lindsleyanum C. B. Adams, 1849, by virtual tautonymy). NOTE: H. B. Baker (1922: 54) found that the radula of the type-species was very close to Lucidella lirata Pfeiffer 1847. Hence, in spite of the Stoastoma-like shell, he placed the taxon tentatively as a subgenus of Lucidella. He also placed Lucidella venezuelensis H. B. Baker 1922, in this subgenus. However, Keen (1960) associated this genus, a synonym of Fadyenia Chitty 1857, with Stoastoma as did Boss (1972).
- lineata C. B. Adams, Helicina, 1845. Proc. Boston Soc. Nat. Hist. 2: 12 (Jamaica; lectotype, MCZ 177223, designated and figured by Jacobson and Boss, 1973, p. 378, pl. 86, figs. 9-11).
- lirata Pfeiffer, Helicina, 1847. Zeits. Malak. 4: 150 (Yucatan, México; type, destroyed); Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2,

p. 341, pl. 68, figs. 5–7. NOTE: lamellosa Guppy 1867, rusticella Morelet 1849, semistriata Sowerby 1866, and unidentata Pfeiffer 1849 are synonyms.

Lucidella Swainson, 1840. Treatise on Malacology, p. 330 (type-species, *Helix (Helicodonta) aureola* Férussac 1822,

by monotypy).

manni Clench and Aguayo, Lucidella, 1932. Proc. New England Zool. Club 13: 35 (Furcy, Haiti, [Hispaniola]; holotype, MCZ 23414); 1937, Mem. Soc. Cubana Nat. His. 11 (2): 63, pl. 7, fig. 8.

midyetti Richards, Lucidella, 1938. Proc. American Philos. Soc. 79: 175, pl. 3, figs. 3-4 (between Coxen Hole and French Harbor, Roatan Id., Honduras; holotype ANSP 170021).

minor "Pfeiffer" Brown, Lucidella aureola, 1913. Proc. Acad. Nat. Sci. Phila., p. 16, nomen nudum. NOTE: we were unable to find this reference in Pfeiffer.

montegoensis Brown, Lucidella aureola, 1913. Proc. Acad. Nat. Sci. Phila. 65: 21, pl. 1, figs. 22–26 (Orange Hill, Montego Bay, Jamaica; type, ANSP 104482a). NOTE: probable synonym of *interrupta* Simpson, 1894, *teste* H. B. Baker (1934).

nana Pfeiffer, Lucidella. 1857. Proc. Zool. Soc. London, p. 111 (Jamaica; syntypes, BMNH 1974.4); Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 342, pl. 68, figs. 12 and 13. NOTE: tridens (Shuttleworth) Sowerby 1866 is a synoym; trochiformis Pilsbry 1899 is probably a synonym (Pilsbry and Brown 1911).

paivana Pfeiffer, Helicina, 1866. Malak. Blätt. 13: 89 (Platon, Haiti, Hispaniola; type, lost); Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 346, pl. 69, figs. 1–2. NOTE: cruciata Weinland 1880 is a synonym, teste Wagner 1910.

peasei Pilsbry, Lucidella (Poeniella), 1933. Proc. Acad. Nat. Sci. Phila. 85: 133, pl. 7, figs 8-8 a (Del Monte) Plantation, 5 or 6 mi W of Barahono, near Salvation, Santa Domingo, 3000 ft., [Hispaniola]; type, ANSP

160986).

- Perenna Guppy, 1867. Ann. Mag. Nat. Hist. (3) 19: 260 (type-species, *Helicina lamellosa* Guppy 1867 [= H. lirata Pfeiffer 1847], by monotypy). NOTE: synonym of *Poenia* H. and A. Adams 1856.
- persculpta Pilsbry and Brown, Lucidella, 1912. Proc. Acad. Nat. Sci. Phila. 63: 584, fig. 1 (Orange Hill and Rosemont Bay, Jamiaca; type, ANSP 104402). NOTE: Baker (1964) gives the type-number as 104415a.
- pilsbryi Clapp, Lucidella, 1914. Nautilus 27: 97, pl. 6, figs. 5 and 6 (Swan Island; [Caribbean Sea]; type, MCZ 22890).
- plicatula Pfeiffer, Helicina, 1849. Proc. Zool. Soc. London,
  p. 123 (Island of Martinique; syntypes, BMNH 1974.2);
  Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab.
  (2) 1: sect. 18, pt. 2, p. 349, pl. 69, figs. 4-7.
- Poenia H. and A. Adams, 1856. Genera of Recent Mollusca 2: 304 (type-species, Helicina depressa Gray 1825, subsequent designation, Pilsbry and Brown 1912). NOTE: H. B. Baker (1922) stated that the type-species was Helicina unidentata Pfeiffer 1849 [= H. lirata Pfeiffer 1847], following the designation by Fischer (1885). However, this designation is invalid because Fisher did not use the word "type." A correct designation was given by Pilsbry and Brown (1912).
- Poeniella H. B. Baker, 1923. Occ. Pap. Mus. Zool., Univ. Michigan, no. 137, p. 23 (type-species, *H. plicatula christophori* Pilsbry 1897, original designation).
- Prosopis Weinland, 1862. Malak. Blätt. 9: 198 (type-species, *Prosopis sulcata* Weinland, 1862, by monotypy), *non* Jurine 1801, *nec* Fabricius 1804 (Hymenoptera). NOTE: synonym of *Lucidella* Swainson 1840.
- rugosa Pfeiffer, Helicina, 1839. Wiegmanns Arch. Naturgesch. 1 (5): 355 (Palenque and Pan de Matanzas, Cuba; type, destroyed); 1848 [in] Martini and Chemnitz, Conch.-Cab. (1) 1: sect. 18, pt. 1, p. 13, pl. 4, figs. 28–30).
- rusticel'a Morelet, Helicina, 1849. Testacea novissima 1: 21 (Carmen Id., Yucatan [Mexico]; syntypes, BMNH 1893.2.4). NOTE: synonym of lirata Pfeiffer 1847.

- samana Pilsbry, Lucidella (Poeniella), 1928. Proc. Acad. Nat. Sci. Phila. 80: 481, pl. 27, figs. 6-7 (1 mi NW of Sanchez, Samana Bay, Santo Domingo, [Hispaniola]; type, ANSP 146696).
- semistriata Sowerby, Helicina, 1866. Thes. Conch. 3: 281, pl. 268 (3), fig. 86 (no locality given; type, destroyed, not in BMNH). NOTE: synonym of *lirata* Pfeiffer 1847.
- striatula Potiez and Michaud, Helicina, 1838. Galérie des Mollusques, Museum de Douai 1: 231, pl. 23, figs. 13–14 (no locality given; type, not located), non Sowerby 1866.
- striatula Sowerby, Helicina, 1842. Proc. Zool. Soc. London, p. 6 (no locality given); Thes. Conch., vol. 1, *Helicina*, p. 3, sp. 6, pl. 1, fig. 43 (Martinique; type BMNH 1974. 3), non Potiez and Michaud 1838.
- striatula (Rang) Pfeiffer, Helicina, 1852. Monographia Pneumonopomorum Viventium, p. 340 (Martinique); "in sched. Cuming," nomen nudum. NOTE: is plicatula Pfeiffer 1849, fide Pfeiffer.
- sublaevis H. B. Baker, Lucidella adamsiana, 1934. Nautilus 48: 9, pl. 2, fig. 4 (nr. Newleigh Hotel, E of Mandeville, S of Williamsfield-Mandeville Road, Jamaica; type, ANSP 162731).
- sulcata Weinland, Prosopis, 1862. Malak. Blätt. 9: 198 (surroundings of Jérémie, Haiti, [Hispaniola]; type, not located).
- tantila (Pilsbry) Vanatta, Lucidella, 1923. Nautilus 37: 69, error for tantilla.
- tantilla Pilsbry, Helicina, 1902. Nautilus 16: 53 (Palm Beach, Florida) holotype, ANSP 77349; 1948. Land Mollusca of North America north of Mexico, Monograph 3, Acad. Nat. Sci. Phila. 2: 1085, fig. 580.
- tridens (Shuttleworth) Sowerby, Lucidella, 1866. Thes. Conch., vol. 3, *Helicina*, p. 282, pl. 268 (3), figs. 90–91 (Jamaica; syntypes, BMNH). NOTE: synonym of *nana* Pfeiffer 1857, *teste* Wagner 1910.
- trochiformis Pilsbry, Lucidella, 1899. Nautilus 13: 56 (Jamaica; type, ANSP 76441a). NOTE: probable synonym of nana Pfeiffer 1857, teste Pilsbry and Brown (1911).

- umbonata Shuttleworth, Helicina, 1854. Mittheil. Naturforsch. Gesell. Bern, nos. 321, 322, p. 93 (Puerto Rico; syntypes, NMB 997); Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 340, pl. 67, figs. 21–24.
- undulata Pfeiffer, Lucidella, 1862. Malak. Blätt. 8: 171 (Jamaica; type, destroyed); Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 338, pl. 67, figs. 16–18.
- unidentata Pfeiffer, Helicina, 1849. Proc. Zool. Soc. London, p. 125 (Honduras; type, probably lost, not in BMNH). NOTE: synonym of lirata Pfeiffer 1847.
- Urichia Guppy, 1895. Proc. Victoria Inst. Trinidad 2: 74 (type-species, *Helicina adamsiana* Pfeiffer 1848, subsequent designation H. B. Baker 1927). NOTE: synonym of *Lucidella* ss.
- valida C. B. Adams, Helicina depressa, 1850. Ann. Lyc. Nat. Hist. New York 5: 52 (Jamaica; lectotype, MCZ 275687, Jacobson and Boss, 1973, Occ. Papers Moll. 3: 437, pl. 86, figs. 5 and 6).
- venezuelensis H. B. Baker, Lucidella (Lindsleya?), 1922. Proc. Acad. Nat. Sci. Phila. 74: 54, nomen nudum; 1923, Occ. Pap. Mus. Zool., Univ. Michigan, no. 134, p. 21, pl. 4, fig. 17 (radula) (near Caño Minapam, Palma Sola, Venezuela; type, ANSP). NOTE: This species and the subgenus Lindsleya, with a shell very much like Stoastoma, were placed by Baker in Lucidella because of striking similarities in the radula with Lucidella lirata Pfeiffer.
- vinosa Shuttleworth, Helicina, 1854. Mittheil. Naturforsch. Gesell. Bern, nos. 321, 322, p. 92 (near San Juan, Humacao, and Rio Blanco, Puerto Rico; type, not found in NMB); Wagner, 1910. [in] Martini and Chemnitz, Conch.-Cab. (2) 1: sect. 18, pt. 2, p. 348, pl. 69, figs. 10 and 11).
- wolffi Boettger, Helicina, 1887. Jahrb. Malak. Gesell. 14: 102, pl. 4, fig. 9 (Miragoane, Haiti, Hispaniola; holotype, SMF 228074).

yallahsensis Pilsbry and Brown, Lucidella 1911. Proc. Acad. Nat. Sci. Phila. 63: 585, fig. 2 (Yallahs, Parish of St. Thomas, Jamaica; type, ANSP 10211).

#### LITERATURE CITED

- ADAMS, C. B. 1851. Description of new species and varieties of the land shells of Jamaica, with notes on some previously described species. Contributions to Conchology, no. 9, pp. 153-174.
- Baker, H. B. 1922. Notes on the radula of *Helicina*. Proc. Acad. Nat. Sci. Phila. 74: 29-67, pls. 3-7.

- Boss, K. J. 1972. Minute Jamaican prosobranch gastropods: Stoastoma and its congeners. Breviora, No. 393, 13 pp., 2 pls.
- ———— and M. K. Jacobson. 1974. Monograph of the genus Lucidella in Cuba (Prosobranchia: Helicinidae). Occ. Papers Moll., Harvard Univ., 4 (48): 1–27, 4 pls.
- Fischer, H. 1885. Helicinidae [in] Manuel de Conchy. pp. 794-798, figs. 554-556.
- HJALMARSON, J. and L. PFEIFFER. 1858. Beiträge zur Fauna von Westindien. Malak. Blatt. 5: 135–155, pls. 2, 3.
- JACOBSON, M. K. 1973. On the identity of *Helicina granulum* (Pfeiffer) (Prosobranchia). Nautilus, 87: 107-109.
- and K. J. Boss. 1973. The Jamaican land shells described by C. B. Adams. Occ. Papers Moll., Harvard Univ., 3(47): 305-519, pls. 54-91.
- KEEN, A. M. 1960. Helicinidae [in] Moore, R. C. (ed.), Treatise on Invertebrate Paleontology, Mollusca, Lawrence, Kansas: Univ. of Kansas Press. Vol. 1, pp. 285–288, figs. 186–187.
- KENNARD, A. S. 1942. The *Histoire* and *Prodrome* of Férussac. Proc. Malac. Soc. London 25: 12-17.
- PILSBRY, H. A. and A. P. BROWN. 1912. The land Mollusca of Montego Bay, Jamaica; with notes on the land Mollusca of the Kingston region. Proc. Acad. Nat. Sci. Phila. (1911) 63: 572-588, figs. 1-2, pl. 43.
- Wagner, A. 1910. Helicinidae [in] Martini and Chemnitz, Conchylien-Cabinet, (2)1: sect. 18, pt. 2, 391 pp., 70 pls.